



Long Range RNA-RNA interactions with the genome of classical swine fever virus

Hadsbjerg, Johanne; Rasmussen, Thomas Bruun; Belsham, Graham

Publication date:
2015

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Hadsbjerg, J., Rasmussen, T. B., & Belsham, G. (2015). *Long Range RNA-RNA interactions with the genome of classical swine fever virus*. Poster session presented at 9th Annual Meeting of EPIZONE, Montpellier, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LONG RANGE RNA-RNA INTERACTIONS

within the genome of classical swine fever virus




Johanne Hadsbjerg, Thomas Bruun Rasmussen, Graham J. Belsham

DTU National Veterinary Institute, Technical University of Denmark, Lindholm, DK-4771 Kalvehave, Denmark

INTRODUCTION

Classical swine fever (CSF) is a highly contagious porcine disease caused by classical swine fever virus (CSFV). The 5'-untranslated region (5' UTR) of the CSFV RNA genome contains an internal ribosomal entry site (IRES) directing the cap-independent initiation of protein synthesis. Besides IRES function, the 5' UTR may also contain *cis*-acting RNA elements that are important not only for viral translation and replication but also important in regulating the switch between these processes.

A motif in the coding region for NS5B (the RNA polymerase) has now been identified in the CSFV RNA which might interact with the IRES subdomain III_{d1}.




NS5B mutants with modifications in a triplet CCC motif

To investigate whether this putative interaction in the CSFV RNA is important in viral replication, a triplet motif of three cytosines (CCC) spanning two codons (xCC | Cxx) within the NS5B coding region was modified to affect the possible interaction with the IRES domain III_{d1} that contains a GGG motif.




METHODS

By site-directed mutagenesis, specific silent mutations have been introduced into the triplet motif within the NS5B sequence and the effect on replication has been analysed using a luciferase reporter system.




RESULTS

All three NS5B mutants have reduced viral replication efficiency. In particular, the mutant with a cytosine to uracil change (resulting in no amino acid change within the protein), exhibited a substantial decrease in replication. This level is comparable to a replication defective mutant denoted mut1.



Replication efficiency in percent compared to wild type (cCc) measured 96 hours post-electroporation



The reduced replication capability indicates that the cCc motif is required for efficient viral replication. This effect is not associated with any change in the amino acid sequence of the RNA polymerase but likely due to interactions within the RNA genome.

Take home messages...

1 A **TRIPLET CCC MOTIF** in the NS5B coding region appears to be important for viral replication

2 **REDUCED REPLICATION EFFICIENCY** is observed after mutating the triplet motif (cAc, cGc, cUc) without changing the encoded protein sequence

3 **UNDERSTANDING PUTATIVE LONG RANGE INTERACTIONS** will provide valuable insight into the mechanisms underlying viral RNA replication

